

SUPERSYMMETRIC WORLDVOLUME SOLITONS

Joan Simón

Departament ECM, Facultat de Física

Universitat de Barcelona and Institut d'Altes Energies

Diagonal 647, E-08028 Barcelona, Spain

jsimon@ecm.ub.es

Abstract The general criteria for finding bosonic supersymmetric worldvolume solitons is reviewed. We concentrate on D-branes, discussing in particular, bion/dyon solutions and D3 branes on NS5 backgrounds.

Introduction. Brane effective actions constitute a relativistic extension of supersymmetric field theories, in the sense that the energy density functional $\mathcal{E}$ satisfies $\mathcal{E}^2 = f(\phi^i)$, which upon linearisation matches the conventional susy free field theory result. This observation suggests that all supersymmetric field theories might be realized in terms of branes and their intersections. It is desirable to establish a complete classification of their classical solutions, especially the BPS ones that will receive no quantum corrections but contribute to the path integral.

A universal feature of all these effective actions is the existence of a fermionic gauge symmetry called kappa symmetry. The latter is not only responsible of removing half of the fermionic degrees of freedom, but it also restricts the background geometry to be compatible with the supergravity equations of motion and relates spacetime supersymmetry with worldvolume supersymmetry. In particular, any bosonic supersymmetric configuration should satisfy [1] :

$$\Gamma_\kappa \epsilon = \epsilon \quad (1.1)$$

where Γ_κ is the matrix appearing in κ -symmetry transformations, satisfying $\Gamma_\kappa^2 = 1$ and $\text{Tr} \Gamma_\kappa = 0$, while ϵ is an arbitrary linear combination of Killing spinors of the background. The number of supersymmetries preserved by the combined background/brane system is the number of

linearly independent solutions of (1.1). Notice that if the background admits no Killing spinors, the worldvolume theory will never be supersymmetric.

It is important to stress that (1.1) is equivalent to the calibration approach [2] for those branes with no excited gauge field living on their worldvolumes. This is the reason why we will concentrate on Dp-branes whose Γ_κ matrix can be written as a $p + 1$ -form (d-form) :

$$\sqrt{-\det(\mathcal{G}_{\mu\nu} + \mathcal{F}_{\mu\nu})} \Gamma_\kappa = e^{\mathcal{F}} \wedge \bigoplus_{n=0}^{(2d-1+(-)^d)/4} \gamma_{2n+\frac{1+(-1)^p}{2}} \mathcal{P}_i^{n+\frac{1+(-1)^p}{2}}$$

where $\gamma_\mu = \partial_\mu X^m e_m^a \Gamma_a$ while $\mathcal{P}_A = \Gamma_{11}$ and $(\mathcal{P}_B)^s = (\sigma_3)^s i\sigma_2$.

Condition (1.1) depends on the background $(e_m^a, B_{mn}, \epsilon)$ and on the configuration (X^m, A_μ) . Once the background is fixed, its solution will not only determine the number of independent components of ϵ but will provide us with an equation(s), that will play the role of BPS equation(s), relating the different excited fields. Once the BPS equation(s) is known, one can always write the square of the energy density as a sum of positive definite terms that allow us to find some bound on the energy [3]. It is precisely when the BPS equation(s) is satisfied that the bound is saturated. Since the energy is minimised, it is no longer necessary to check the equations of motion, but the Gauss' law and/or the Bianchi identity, that will determine the equation of motion for the remaining independent excited fields.

Bions. Let us consider Dp-branes probing Dp-branes geometry ($1 \leq p \leq 7$). The background is given by

$$ds^2 = U^{-1/2} ds^2(\mathbb{M}_{(1,p)}) + U^{-1/2} ds^2(\mathbb{E}_{9-p}) \quad (1.2)$$

$$e^\phi = U^{(3-p)/4} \quad C_{0\dots p} = U^{-1} \quad (1.3)$$

where the gauge choice for the R-R potential is consistent with the no force condition existing between parallel Dp-branes and U is an harmonic function in the transverse space to the branes.

To describe a F-string ending on a probe parallel to a stack of Dp-branes $(X^\mu = \xi^\mu, \mu = 0, \dots, p)$, we must excite a transverse scalar $(Y = Y(\xi^a), a = 1, \dots, p)$ and the electric components of the gauge field $A_0 = A_0(\xi^a)$, since the F-string is an electrically charged NS-NS object. It is natural to demand $\Gamma_{0\dots p} \mathcal{P}_i^{(2d+1-(-)^d)/4} \epsilon_0 = \epsilon_0$ and $\Gamma_{0y} \Gamma_{11} \epsilon_0 = -\epsilon_0$ for type IIA and $\Gamma_{0y} \sigma_3 \epsilon_0 = \epsilon_0$ for type IIB, on the asymptotic value of the Killing spinor (ϵ_0), since these are the conditions describing the existence of a Dp-brane and a F-string, respectively.

When eq. (1.1) is subject to the above constraints, it is solved iff

$$F_{0a} = \partial_a Y \quad \Leftrightarrow \quad \tilde{E}^a = \delta^{ab} \partial_b Y \quad (1.4)$$

where $\tilde{E}^a$ is the conjugate momentum of the gauge field. (1.4) is the BPS equation as it can be checked by analysing the energy density [4] of the configuration

$$(\mathcal{E} + U^{-1})^2 = (U^{-1} + \tilde{E}^a \partial_a Y)^2 + U^{-1} |\tilde{E}^a - \delta^{ab} \partial_b Y|^2. \quad (1.5)$$

The Gauss' law ($\partial_a \tilde{E}^a = 0$) determines the eq. of motion ($\delta^{ab} \partial_a \partial_b Y = 0$) for the excited scalar. Its solution and the corresponding physical interpretation depend on the worldvolume dimension.

Due to the self-duality of $G_{(5)} = dC_{(4)}$, one should expect dyonic solutions in the case of D3-branes. The soliton condition on ϵ_0 is modified to $\Gamma_{0y}(\cos \alpha \sigma_3 + \sin \alpha \sigma_1) \epsilon_0 = \epsilon_0$, while (1.1) is solved iff $F_{0a} = \cos \alpha \partial_a Y$ and $B^a = \sin \alpha \delta^{ab} \partial_b Y$. The excitation of the magnetic field is consistent with the charge carried by the D-strings.

The latter analysis could be extended to D8-branes, since there is no new contribution due to the mass parameter for the configurations under study.

D3-branes in NS5 backgrounds. Let us consider a D3-brane probing NS5-branes geometry. The background is given by :

$$ds^2 = ds^2(\mathbb{M}_{1,5}) + U \left(dr^2 + r^2 d\Omega_3^2 \right) \quad (1.6)$$

$$e^{2\phi} = U, \quad H_{(3)} = 2R^2 \omega_{(3)} \quad (1.7)$$

We will describe the S-dual configuration to the one considered in [5] (to which we refer for notation), but from the D3-brane perspective. The D3-brane "wraps" the transverse S^3 sphere and the NS5-branes magnetic charge induces a magnetic charge on the probe through a D-string soliton connecting both. In this way, the asymptotic Killing spinor ϵ_0 should satisfy $\Gamma_{0\dots 5} \sigma_3 \epsilon_0 = \epsilon_0$, $\Gamma_{0\theta\theta_2\theta_3} i\sigma_2 \epsilon_0 = \epsilon_0$ and $\Gamma_{0r} \sigma_1 \epsilon_0 = -\epsilon_0$, that correspond to the background, probe and soliton susy projection conditions.

Due to the polar coordinates on $\mathbb{E}^4$, ϵ is not constant. Proceeding as in [5], eq. (1.1) is solved if :

$$\begin{aligned} B^{\hat{a}} &= -\Delta \frac{\bar{g}^{\hat{a}\hat{b}} \partial_{\hat{b}} r}{r} \\ B^{\theta} &= \frac{-\Delta}{(r \sin \theta)'} \left((r \cos \theta)' - \bar{g}^{\hat{a}\hat{b}} \frac{\partial_{\hat{a}} r \partial_{\hat{b}} (r \sin \theta)}{r} \right) \end{aligned} \quad (1.8)$$

where $\Delta = U r^2 \sqrt{\det \bar{g}}$ and $B^a = \frac{1}{2} \epsilon^{abc} \mathcal{F}_{bc}$. Eqs. (1.8) are the BPS equations as can be checked by writting the energy density as a sum of squares :

$$\begin{aligned} \mathcal{E}^2 &= |r B^{\hat{a}} + \Delta \bar{g}^{\hat{a}\hat{b}} \partial_{\hat{b}} r|^2 + (B^a \partial_a (r \sin \theta) + \Delta (r \cos \theta)')^2 + \mathcal{Z}^2 \\ \mathcal{Z} &= \Delta (r \sin \theta)' - B^a \partial_a (r \cos \theta) \end{aligned} \quad (1.10)$$

The Bianchi identity $d\mathcal{F} = -d\hat{B}$, which is equivalent to $\partial_a B^a = 2R^2 \sqrt{\det \bar{g}}$, not only gives the corresponding equation of motion for the excited radial scalar, but it is also useful to check the topological character of $\mathcal{Z}$ which can be written as $\mathcal{Z} = \partial_a \mathcal{Z}^a$, where

$$\mathcal{Z}^{\hat{a}} = -B^{\hat{a}} r \cos \theta \quad (1.11)$$

$$\mathcal{Z}^{\theta} = -B^{\theta} r \cos \theta + \sqrt{\det \bar{g}} \sin \theta \left(\frac{r^3}{3} + r R^2 \right) \quad (1.12)$$

A similar analysis to the one carried in [5] shows that for $SO(2)$ invariant configurations, the above soliton gives a worldvolume realization of the Hannany-Witten effect.

Acknowledgments

J. S. would like to thank the organizers for a very nice school, and for financial support. This work was supported by a fellowship from Comissionat per a Universitats i Recerca de la Generalitat de Catalunya and by AEN98-0431 (CICYT), GC 1998SGR (CIRIT).

References

- [1] E. Bergshoeff, R. Kallosh, T. Ortín and G. Papadopoulos, *κ -symmetry, supersymmetry and intersecting branes*, Nucl. Phys. **B502** (1997) 149.
- [2] J. Gutowski, G. Papadopoulos and P. K. Townsend, *Supersymmetry and Generalized Calibrations*, hep-th/9905156.
- [3] J. P. Gauntlett, J. Gomis and P. K. Townsend, *BPS bounds for worldvolume branes*, JHEP 9801:003.
- [4] E. Bergshoeff and P. K. Townsend, *Super D-branes revisited*, Nucl. Phys. **B531** (1998) 226.
- [5] J. Gomis, A. V. Ramallo, J. Simón and P. K. Townsend, *Supersymmetric baryonic branes*, hep-th/9907022.